

Tier 1 Examination - Algebra

January 4, 2001

Justify all answers! All rings are assumed to have an identity element. The set of real numbers is denoted by \mathbf{R} . The set of integers modulo n is denoted \mathbf{Z}_n . The group of permutations on n letters is denoted S_n .

- (20)1. Give an example of each of the following. No justification is required.
- (a) A nonabelian group of order 18.
 - (b) An infinite commutative ring R such that for all $y \in R$, $y + y + y = 0$.
 - (c) A 3 by 3 real matrix that is diagonalizable over the complex numbers but not over the reals.
 - (d) A unique factorization domain that is not a principal ideal domain.
 - (e) An element of order 3 in $\mathbf{GL}_2(\mathbf{R})$.
- (10) 2. Let G be a group with the property that $g^2 = e$ for all $g \in G$. Prove G is abelian.
- (10)3. Determine the number of homomorphisms from S_3 to $\mathbf{Z}_2 \times \mathbf{Z}_4$.
- (10)4. Find $\lim_{n \rightarrow \infty} \begin{pmatrix} 2 & 3 \\ -1/2 & -1/2 \end{pmatrix}^n$.
- (10)5. Let n be a positive odd integer and let $A, B \in M_n(\mathbf{R})$ such that $A^2 = B^2 = I$. Prove that A and B have a common eigenvector (not necessarily with the same eigenvalue).
- (10)6. Let R and S be commutative rings and let $\phi : R \rightarrow S$ be a ring homomorphism. Suppose there is an ideal I of R such that $\ker(\phi) \subset I \subset R$ (proper containments). Prove that the image of ϕ is not a field.
- (10)7. Let F be a subfield of \mathbf{R} and suppose m and n are positive integers with $\sqrt{m} + \sqrt{n} \in F$. Prove that \sqrt{m} and \sqrt{n} are in F .
- (10)8. Let K be a field and let K^\times denote the group of nonzero elements of K . Prove that K^\times contains at most two elements of order 6.

(10)9. Let $G = (\mathbb{Q}, +)/(\mathbb{Z}, +)$, where $(\mathbb{Q}, +)$ denotes the group of rational numbers under addition and $(\mathbb{Z}, +)$ denotes the subgroup of integers under addition. Prove that G is an infinite group in which every element has finite order.

(15)10. (a) Let R be a commutative ring and suppose I and J are ideals of R such that $I + J = R$, where $I + J = \{i + j \mid i \in I, j \in J\}$. Prove the map $\phi : R/I \times R/J \rightarrow R/(I \cap J)$ given by $\phi(r + I, r + J) = (r + I \cap J)$ is an isomorphism of rings.

(b) Let R be a commutative ring containing exactly 4 ideals (including $\{0\}$ and R). Let I and J denote the other two ideals and suppose they are incomparable, that is $I \not\subseteq J$ and $J \not\subseteq I$. Prove that R is isomorphic to the direct product of two fields.