

# Tier 1 Analysis Examination

August 1997

1. Does  $a_k = \frac{1 \cdot 3 \cdot 5 \cdot 7 \cdot 9 \cdots (2k+1)}{2 \cdot 4 \cdot 6 \cdots 2k}$  converge or diverge? Prove your assertion.

2. Let  $S \subset \mathbb{R}^3$  be the "tin can without a lid".

$$\{(x, y, z) : x^2 + y^2 = 1, 0 \leq z \leq 1\} \cup \{(x, y, z) : x^2 + y^2 \leq 1, z = 0\}$$

Compute the flow  $\iint_S F \cdot N \, dA$  "out" of the can if  $F = (x(y+z), -zy, -zy)$

3. Definition: A transformation of class  $C^1$   $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is called volume preserving if for every cube  $C \subset \mathbb{R}^3$ , with faces parallel to the coordinate planes,  $\text{volume}(F(C)) = \text{volume}(C)$ .

(i) Show that  $F(x, y, z) = (x + y, z - 4, z^2 - y)$  is volume preserving.

(ii) Show that if  $G : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  is volume preserving then the determinant of its derivative  $G'$  equals  $\pm 1$ , and  $G$  maps open sets into open sets.

4. Let  $f_n(x) = \int_{1/2}^x \arctan^2(t/n) \, dt \quad n = 1, 2, \dots$

(i) Show that  $\sum_{n=1}^{\infty} f'_n(x)$  (sum of derivatives) is uniformly convergent on  $[-1, 1]$ .

(ii) Show that  $g(x) = \sum_{n=1}^{\infty} f_n(x)$  is differentiable for all  $x$ .

5. Let  $I_j$  be a countable family of closed intervals whose interiors are pairwise disjoint and such that  $\bigcup I_j = [0, 1]$ . Show directly (without fancy integration theorems) that  $\sum_{j=1}^{\infty} |I_j| = 1$ .

6. Give a counterexample to this statement: Every  $f : \mathbb{R} \rightarrow \mathbb{R}$  with the property that  $f^{-1}(K)$  is compact for any compact  $K$  is continuous.

7. Let  $g : \mathbb{R}^2 \rightarrow \mathbb{R}$  be a  $C^\infty$  function with  $0 \in \mathbb{R}^2$  a critical point. Suppose the matrix of second partials  $\left(\frac{\partial^2 g}{\partial x_i \partial x_j}(0)\right)$  has eigenvalues  $-2$  and  $0$ . Show that the origin is NOT a local minimum of  $g$ .

8. Definition: A metric space is said to have property  $Z$  if every sequence with exactly one cluster point converges. (Recall that every neighborhood of a cluster point of  $\{x_n\}$  contains infinitely many  $x_n$ ).

(i) Give an example of a metric space that has property  $Z$  and an example of a metric space that does not.

(ii) What properties of metric spaces are implied by or equivalent to property  $Z$ ?