

Tier I Analysis Exam
August, 2005

Justify your answers. All problems carry equal weight.

1. Let $p > 0$. Evaluate

$$\lim_{n \rightarrow \infty} \frac{1^p + 3^p + \dots + (2n-1)^p}{n^{p+1}}$$

2. For $x > 0$, define

$$\phi(x) = \begin{cases} 0 & \text{if } x \text{ is irrational} \\ \frac{1}{q} & \text{if } x = \frac{p}{q} \end{cases}$$

where p, q have no common factor and $q \geq 1$.

- (a) Where is ϕ continuous?
(b) Where is ϕ differentiable?

3. Let a, b be real with $|b| > \max\{1, |a|\}$. For $x \in \mathbf{R}$, define

$$f(x) = \sum_{n=1}^{\infty} \frac{\cos(a^n x)}{b^n}.$$

- (a) Show that f is uniformly continuous on all of \mathbf{R} .
(b) Let

$$\gamma = \{(x, y) : y = f(x), 0 \leq x \leq 1\}$$

be the graph of f over the unit interval. Show that γ has finite length.

4. Let M_2 denote the set of 2-by-2 matrices with real entries, and for $A \in M_2$, define $S(A) = A^2$. Does the mapping $S : M_2 \rightarrow M_2$ have a local inverse near the identity matrix?
5. Fix $a > 0$. Let x_1, \dots, x_n be non-negative numbers with

$$\sum_{i=1}^n x_i = na.$$

Show that

$$\sum_{i < j} x_i x_j \leq \frac{1}{2} n(n-1) a^2.$$

Please turn over.

6. Let p be real. Suppose $f : \mathbf{R}^n - \{0\} \rightarrow \mathbf{R}$ is continuously differentiable, and satisfies

$$f(\lambda x) = \lambda^p f(x), \text{ for all } x \neq 0 \text{ and for all } \lambda > 0.$$

Let $\nabla f(x)$ denote the gradient of f at x and \cdot the Euclidean inner product. Prove that

$$x \cdot \nabla f(x) = p f(x), \quad x \neq 0.$$

7. A family \mathcal{F} of continuous real-valued functions of a real variable is called *equicontinuous at x* if for every $\epsilon > 0$, there is a $\delta > 0$ such that for every $f \in \mathcal{F}$,

$$|x - y| < \delta \Rightarrow |f(x) - f(y)| < \epsilon.$$

\mathcal{F} is called *equicontinuous on the set E* if it is equicontinuous at each point x of E . (Note: the constant δ may depend on both ϵ and x .)

Now suppose \mathcal{F} is a family of continuous real-valued functions defined on an open interval $I \subseteq \mathbf{R}$, and let $x_0 \in I$.

- (a) Suppose \mathcal{F} is equicontinuous at every point of $I \setminus \{x_0\}$. Must \mathcal{F} also be equicontinuous at x_0 ?
- (b) Suppose \mathcal{F} is equicontinuous at x_0 . Must \mathcal{F} also be equicontinuous at every point in some neighborhood J of x_0 ?
8. Let U be an open subset of \mathbf{R}^n and $f : U \rightarrow \mathbf{R}^n$ be differentiable. Suppose there exists $C > 0$ such that

$$|f(x) - f(y)| \geq C|x - y|$$

for all $x, y \in U$. Let $df(x)$ denote the Jacobian derivative of f at x (that is, the linear mapping given by the n by n matrix of partial derivatives). Show that $\det df(x) \neq 0$ for all $x \in U$.

9. Let $m > 0$ be a real number, let $r = (x^2 + y^2 + z^2)^{1/2}$, and consider the vector field on \mathbf{R}^3 given by $\vec{F} = r^m \cdot (x, y, z) = (x^2 + y^2 + z^2)^{m/2}(x, y, z)$.
- (a) Compute the divergence $\text{div}(\vec{F})$.
- (b) Using part (a) and the Divergence Theorem, calculate

$$\iiint_{B^3} r^m dV$$

where $B^3 = \{(x, y, z) : r \leq 1\}$ is the closed unit ball centered at the origin and $dV = dx dy dz$ is the Euclidean volume.

End of exam.