

**TIER 1 ANALYSIS EXAM
AUGUST 2007**

- (1) Define $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ by setting

$$f(x, y) = \frac{x^3 + y^3}{x^2 + y^2}$$

for $(x, y) \neq (0, 0)$ and $f(0, 0) = 0$. Show that f is differentiable at all points $(x, y) \in \mathbb{R}^2$ except $(0, 0)$. Show that f is not differentiable at $(0, 0)$.

- (2) Given $\lambda \in \mathbb{R}$, define $h_\lambda : \mathbb{R}^2 \rightarrow \mathbb{R}$ by

$$h_\lambda(x, y) = -x^4 + x^2 + y^2 + \lambda \cdot \sin(x \cdot y).$$

For which values of λ does h_λ have a local minimum at $(0, 0)$? Justify your answer.

- (3) Let $\gamma \subset \mathbb{R}^2$ be the simple closed curve described in polar coordinates by $r = \cos(2\theta)$ where $\theta \in [-\pi/4, \pi/4]$. Suppose that γ is positively oriented. Compute the line integral

$$\int_{\gamma} 3y \, dx + x \, dy.$$

Provide the details of your computation.

- (4) Let X be a metric space such that $d(x, y) \leq 1$ for every $x, y \in X$, and let $f : X \rightarrow \mathbb{R}$ be a uniformly continuous function. Does it follow that f must be bounded? Justify your answer with either a proof or a counterexample.

- (5) Let

$$f(x, y) = (x + e^{2y} - 1, \sin(x^2 + y)),$$

and let

$$h(x, y) = (1 + x)^5 - e^{4y}.$$

Show that there exists a continuously differentiable function $g(x, y)$ defined in a neighborhood of $(0, 0)$ such that $g(0, 0) = 0$ and $g \circ f = h$. Compute $\frac{\partial g}{\partial y}(0, 0)$.

- (6) Let c_1, c_2, \dots be an infinite sequence of distinct points in the interval $[0, 1]$. Define $f : [0, 1] \rightarrow \mathbb{R}$ by setting $f(x) = 1/n$ if $x = c_n$ and $f(x) = 0$ if $x \notin \{c_n\}$. State the definition of a Riemann integrable function, and directly use this definition to show that

$$\int_0^1 f(x) dx$$

exists.

- (7) Show that the formula

$$g(x) = \sum_{n=1}^{\infty} \frac{1}{n^2} e^{\int_0^x t \sin(\frac{n}{t}) dt}$$

defines a function $g : \mathbb{R} \rightarrow \mathbb{R}$. Prove that g is continuously differentiable.

- (8) Consider an unbounded sequence $0 < a_1 < a_2 < \dots$, and set

$$s = \limsup_{n \rightarrow \infty} \frac{\log n}{\log a_n}.$$

Show that the series

$$\sum_{n=1}^{\infty} a_n^{-t}$$

converges for $t > s$ and diverges for $t < s$.

- (9) Define a sequence $\{a_n\}$ by setting $a_1 = 1/2$ and $a_{n+1} = \sqrt{1 - a_n}$ for $n \geq 2$. Does the sequence a_n converge? If so, what is the limit? Justify your answer with a proof.