

# Tier 1 Analysis Exam

January 2021

Work all nine problems. They all count equally. Show computations and justify your answers; a correct answer without a correct proof earns little credit. Write a solution of each problem on a separate page. You have 4 hours.

**Notation:** For a function  $f: \mathbb{R}^n \rightarrow \mathbb{R}$ , denote the partial derivative in the  $i$ -th coordinate direction by  $D_i f$ . The partial derivative of  $D_i f$  in the  $j$ -th coordinate direction is likewise denoted by  $D_{ij} f := D_j D_i f$ . The expression  $:=$  is used to indicate a definition.

1. Let  $B(K)$  denote the set of bounded functions  $f: K \rightarrow \mathbb{R}$ , where  $K \subset \mathbb{R}$  is compact.

(a) For  $f, g \in B(K)$ , define

$$d(f, g) := \sup_{x \in K} |f(x) - g(x)|.$$

Show that  $d: B(K) \times B(K) \rightarrow \mathbb{R}$  defines a metric on  $B(K)$ .

(b) Show that the set  $C(K)$  of continuous functions  $f: K \rightarrow \mathbb{R}$  is a closed subset of  $B(K)$  in the topology given by this metric.

2. Let  $a_{n,m} \in [0, 1]$  for all positive integers  $n$  and  $m$ . Suppose that for each  $n$ , we have  $\lim_{m \rightarrow \infty} a_{n,m} = n/2^n$ . For each of the following inequalities, prove that it must hold or prove (with a counterexample) that it need not hold:

(a)  $\liminf_{m \rightarrow \infty} \sum_{n=1}^{\infty} \frac{a_{n,m}}{n} \geq 1$ ;

(b)  $\limsup_{m \rightarrow \infty} \sum_{n=1}^{\infty} \frac{a_{n,m}}{n} \leq 1$ .

3. Let  $a_k \geq 0$  for all nonnegative integers  $k$ . Suppose that  $\sum_{k=0}^{\infty} a_k < \infty$ . Define  $f(x) := \sum_{k=0}^{\infty} a_k x^k$  for  $x \in [0, 1]$ . Do not assume that  $b := \sum_{k=0}^{\infty} k a_k$  is finite.

(a) Show that (the left-hand derivative)  $f'(1) = b$ .

(b) Show that  $\lim_{x \rightarrow 1^-} f'(x) = b$ .

4. Let  $a_k \geq 0$  for all nonnegative integers  $k$ . Suppose that  $\sum_{k=0}^{\infty} a_k = 1$ ,  $\sum_{k=0}^{\infty} k a_k = 1$ , and  $c := \sum_{k=0}^{\infty} k(k-1) a_k$  is finite. Define  $f(x) := \sum_{k=0}^{\infty} a_k x^k$  for  $x \in [0, 1]$ . You may assume without proof that  $\lim_{x \rightarrow 1^-} f'(x) = 1$  and  $\lim_{x \rightarrow 1^-} f''(x) = c$ . (These both follow from problem 3(b).) Define

$$g(x) := \frac{1}{1 - f(x)} - \frac{1}{1 - x}$$

for  $x \in [0, 1)$ . Show that  $\lim_{x \rightarrow 1^-} g(x) = c/2$ .

5. Consider a function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ . Suppose that  $D_1f$  exists at  $(0, 0)$ . Suppose also that  $D_2f$  exists in a neighborhood of  $(0, 0)$  and is continuous at  $(0, 0)$ . Prove that  $f$  is differentiable at  $(0, 0)$ .

6. Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be twice continuously differentiable in a neighborhood of  $(0, 0)$ , with  $D_2f(0, 0) = 0$  and  $D_{22}f(0, 0) > 0$ .

(a) Prove that there are  $\epsilon, \delta > 0$  such that for each  $x \in (-\epsilon, \epsilon)$ , the formula

$$g(x) := \min\{f(x, y) : |y| \leq \delta\}$$

defines a differentiable function  $g: (-\epsilon, \epsilon) \rightarrow \mathbb{R}$ .

(b) Find a formula for  $g'(0)$  in terms of  $f$  and prove it.

7. Consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  given by  $f(x, y) := x^4 + y^4 - 4xy$ .

(a) Identify and classify all critical points of  $f$  on  $\mathbb{R}^2$ .

(b) Determine the minimum and maximum values of  $f$  on the curve  $x^4 + y^4 = 32$ .

8. Let  $\mathbb{Q} \subset \mathbb{R}$  denote the rationals. Show that the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  defined by

$$f(x, y) := \begin{cases} x & \text{if } (x, y) \in \mathbb{Q} \times \mathbb{Q}, \\ y & \text{otherwise} \end{cases}$$

is not Riemann integrable on the unit square  $[0, 1] \times [0, 1] \subset \mathbb{R}^2$ .

9. Let  $S := \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 + z^2 = 1 \text{ with } y \geq 0 \text{ and } z \geq 0\}$  be the surface consisting of a quarter of the unit sphere in  $\mathbb{R}^3$ . Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be infinitely differentiable. Define the vector field  $\mathbf{F}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  by  $\mathbf{F}(x, y, z) := (f(z) + x^2y, xy^2 + 1, xyz)$ . Evaluate the surface integral  $\int_S \mathbf{F} \cdot \hat{n} \, dS$ , where  $\hat{n}$  is the unit normal vector field of  $S$  pointing away from the origin.